

Black Spot of Rose (*Diplocarpon rosea*)
 Botrytis (*Botrytis* spp.)
 Downy Mildew (*Bremia lactucae*, *Peronospora* spp. and *Plasmopara viticola*) – suppression
 Early Blight (*Alternaria solani*) – suppression
 Fire Blight (*Erwinia amylovora*) – suppression
 Gray Mold (*Botrytis cinerea*)
 Late Blight (*Phytophthora infestans*) – suppression
 Leaf Spots (*Alternaria* spp., *Cercospora* spp., *Entomospodium* spp., *Helminthosporium* spp., *Myrothecium* spp., *Septoria* spp.)
 Pin Rot (*Alternaria/Xanthomonas* complex) – suppression
 Powdery Mildew (*Uncinula necator*, *Erysiphe* spp., *Sphaerotheca* spp., *Oidiopsis taurica*, *Leveillula taurica*, *Podosphaera leucotricha*, *Oidium* spp., *Podosphaera* spp.)
 Scab (*Venturia* spp.) – suppression
 Walnut Blight (*Xanthomonas campestris*)

STORAGE AND DISPOSAL

STORAGE: Store in original container only, in an area inaccessible to children. Keep container closed when not in use.
CONTAINER DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available.
 If partly filled: Call your local waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.
 For emergencies such as leaks or spills contact CHEMTREC 24-hour toll-free hotline at 1-800-424-9300.

Questions or Comments call 800-728-0014
 AgraQuest, Inc., 1540 Drew Avenue, Davis, CA 95618
 www.agraquest.com

CONDITIONS FOR SALE AND WARRANTY

Except to the extent prohibited by applicable law, AgraQuest offers this product with the following conditions: 1) buyers and users of this product assume the risk of any storage, handling or use contrary to AgraQuest's label and directions and 2) AgraQuest's liability shall in no case exceed the purchase price of the applicable AgraQuest product.

U.S. Patent Nos. 6,060,051, 6,103,228, 6,291,426, and 6,417,163 on QST 713 strain of *Bacillus subtilis*
 Serenade® is a registered trademark of AgraQuest, Inc.
 © Copyright AgraQuest, Inc., 2007



OMRI™
Listed
 Organic Materials Review Institute



Serenade®
 — garden —

DISEASE CONTROL
 — READY TO USE —

Controls Bacterial Spot, Powdery Mildew, Rust, Gray Mold, Leaf Blight, Scab

AgraQuest, Inc., 1540 Drew Ave., Davis, CA 95618

Questions or Comments
 call 800-728-0014
 www.agraquest.com

EPA Reg. No. 69592-16
 EPA Est. No. 69592-MEX-1

Serenade® is a registered trademark of AgraQuest, Inc.
 Copyright AgraQuest, Inc. 2007
 Made in Mexico
 73XB-001
 US0005-B-002



FF-8931

**PRECAUTIONARY
STATEMENTS**

**HAZARDS TO HUMANS &
DOMESTIC ANIMALS
CAUTION**

Harmful if inhaled. Avoid breathing spray mist.
Remove contaminated clothing and wash before
reuse.

As a general precaution, when exposed to a living microbial
product such as this, wear a dust particle mask when applying
this product.

FIRST AID

IF INHALED: Move person to fresh air. If person is not breathing,
call 911 or an ambulance, then give artificial respiration,
preferably mouth-to-mouth if possible. Call a doctor or poison
control center for further treatment advice. Have the product
label with you when calling a doctor or poison control center.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when
disposing of equipment wash waters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner
inconsistent with its labeling.

GENERAL USE INFORMATION

Serenade Garden Disease Control Ready To Use (RTU) is a broad
spectrum, preventative biofungicide recommended for the control
or suppression of many important plant diseases and may be
used on roses, vegetables, fruits, nuts, flowers, houseplants,
foliage, trees, shrubs located in residential landscapes, residential
greenhouses, commercial landscapes, and commercial interior
scapes.

APPLICATION INSTRUCTIONS

Shake well before using. Spray leaves, stems and new shoots to
runoff providing complete coverage of the entire plant. For best
results, treat prior to foliar disease development or at the first sign
of foliar disease infection. Repeat at 7-day intervals or as
needed. Under conditions of high disease pressure, when
environmental conditions favor rapid disease development (high
humidity, excessive rain, extreme moisture condition, etc.) shorten
the spray interval. Do not water foliage within 4 hours of
application.

Serenade Garden Disease Control Ready To Use can be applied
up to and including the day of harvest.

**SERENADE
GARDEN DISEASE
CONTROL READY TO
USE MAY BE USED ON THE
FOLLOWING:**

HOME and GARDEN VEGETABLES,
FRUITS AND NUTS: Broccoli, Onion, Melon,
Pepper, Tomato, Lettuce, Apple, Pear, Carrot, Walnut
GREENHOUSE PLANTS: Broccoli, Onion, Melon,
Pepper, Tomato, Lettuce, Carrot
ORNAMENTALS, TREES, SHRUBS, FLOWERING PLANTS,
TROPICAL PLANTS:

It is impossible to test all plants for phytotoxicity. To assure that
the plants to be treated are not sensitive to the treatment, apply
a small amount of the highest application rate of the product to
a few leaves or the above ground portion of a plant and check
within 3 days. Use product according to label directions.

PLANTS EVALUATED FOR PHYTOTOXICITY

Annual and Perennial Flowering Plants: Alyssum, Asters,
Azalea, Begonia, Calla lily, Chrysanthemum, Cyclamen,
Dianthus, Dwarf Bee-Balm, Easter lily, Garden phlox,
Geraniums, Gerbera, Golden star, Hydrangea, Impatiens,
Kalanchoe, Linaria, Lisianthus, Lobelia, Marigolds, Orchids,
Pansies, Petunia, Poinsettia, Portulaca, Ranunculus, Roses,
Salvia spp., Snapdragons, Stock, *Verbena* spp., Vinca, Violas,
Zinnias

Tropical foliage: Aglaonema, Dieffenbachia, *Dracaena* spp.,
English Ivy, Hibiscus, Leatherleaf Fern, Spathiphyllum

Trees and Shrubs: Azalea, Boxwood, Crape myrtle, Dogwood,
Gumbo azalea, Indian Hawthorn, Japanese maple, *Ligustrum
japonicum*, Lilac, Loropetalum, Photinia, Rhododendron,
Rosaceae spp., Soft Touch Holly, Spirea.

**DISEASES CONTROLLED OR SUPRESSED BY
SERENADE GARDEN DISEASE CONTROL READY
TO USE ON VEGETABLES, FRUIT, NUTS,
ORNAMENTAL PLANTS**

Anthraxnose (*Colletotrichum* spp.)
Bacteria (*Erwinia* spp., *Psuedomonas*, *Xanthomonas* spp.)
Bacterial Leaf Blight (*Xanthomonas campestris*)
Bacterial Speck (*Psuedomonas syringae* pv. Tomato)
Bacterial Spot (*Xanthomonas* spp.) – suppression
Black Mold (*Alternaria alternata*)
Black Root Rot/Black Crown Rot (*Alternaria* spp.)